

# A Hyperexponential Approximation to Finite- and Infinite-time Ruin Probabilities of Compound Poisson Processes

Amir T. Payandeh Najafabadi, Member IAENG \*

*Abstract*—This article considers the problem of evaluating infinite-time (or finite-time) ruin probability under a given compound Poisson surplus process. By approximating the claim size distribution by a finite mixture exponential, say Hyperexponential, distribution. It restates the infinite-time (or finite-time) ruin probability as a solvable ordinary differential equation (or a partial differential equation). Application of our findings has been given through a simulation study.

*Keywords:* Ruin probability, Compound Poisson Processes, mixture exponential (Hyperexponential) distribution, Heavy-tailed distributions

## 1 Introduction

Consider the following compound Poisson process

$$U_t = u + ct - \sum_{j=1}^{N(t)} X_j, \quad (1)$$

where  $X_1, X_2, \dots$  are a sequence of i.i.d. random variables with common density function  $f_X(\cdot)$ ,  $N(t)$  is a Poisson process with intensity rate  $\lambda$ ,  $u$  and  $c$  stand for initial wealth/reserve and premium of the process, respectively.

The finite-time and infinite-time ruin probabilities for the above compound Poisson process are, respectively, denoted by  $\psi(u; T)$  and  $\psi(u)$  and defined by

$$\psi(u; T) = P(\tau_u \leq T); \quad \& \quad \psi(u) = P(\tau_u < \infty), \quad (2)$$

where  $\tau_u = \inf\{t : U_t \leq 0 | U_0 = u\}$  is the hitting time.

Ref. [15] among others, established that an infinite-time ruin probability  $\psi(u)$  under a compound Poisson process can be restated as the following integro-differential equation

$$c\tilde{\psi}^{(1)}(u) - \lambda\tilde{\psi}(u) + \lambda \int_0^u \tilde{\psi}(u-x)f_X(x)dx = 0, \quad (3)$$

where  $\tilde{\psi}(u) = 1 - \psi(u)$  and  $\lim_{u \rightarrow \infty} \psi(u) = 0$ .

A finite-time ruin probability  $\psi(u; T)$  under a compound Poisson process can be restated as the following partial integro-differential equation

$$\begin{aligned} \frac{\partial \tilde{\psi}(u; T)}{\partial u} - \frac{\partial \tilde{\psi}(u; T)}{\partial T} - \frac{\lambda}{c} \tilde{\psi}(u; T) \\ + \frac{\lambda}{c} \int_0^u \tilde{\psi}(u-x; T) f_X(x) dx = 0, \end{aligned} \quad (4)$$

where  $\tilde{\psi}(u; T) = 1 - \psi(u; T)$ ,  $\lim_{u \rightarrow \infty} \psi(u; T) = 0$  for all  $T > 0$  and  $\psi(u; T = 0) = 0$  for all  $u \geq 0$ , see Ref [23].

Since the compound Poisson surplus process plays a vital role in many actuarial models, several authors studies ruin probability under surplus process (1). An excellent review for infinite-time ruin probability can be found in [3]. For finite-time ruin probability: [23] showed for exponential claim size distribution partial integro-differential equation (4) can be transformed into a second-order partial differential equation. Ref. [1] considered a compound Poisson surplus process with constant force of real interest. Then, they restated finite-time ruin probability  $\psi_T(u)$  as a gamma series expansion. Ref. [14] provided a global Lagrange type approximation in the z-space for  $\psi_T(u)$  under surplus process (1). Ref. [2] and [29] employed the Padé approximant method to approximate  $\psi_T(u)$  under surplus process (1).

This article in the first step approximates claim size density function  $f_X(\cdot)$  with a finite mixture exponential, say Hyperexponential, density function  $f_X^*(\cdot)$ . Then, it transforms two integro-differential equations (3) and (4), respectively, into an ordinary differential equation (ODE) and a partial differential equation (PDE). A simulation study has been conducted to show practical application of our findings.

The rest of this article is organized as follows. Some mathematical background for the problem has been collected in Section 2. Section 3 provides the main contribution of this article. Applications of the results have been given in Section 4.

## 2 Preliminaries

From hereafter now, we set  $\sum_{j=a}^b A_j = 0$ , for  $b < a$ .

\*A. T. Payandeh Najafabadi is with the Department of Mathematical Sciences, Shahid Beheshti University, G.C. Evin, 1983963113, Tehran, Iran. e-mail: amirtpayandeh@sbu.ac.ir.



tribution; (2) Error bound for such approximation satisfies  $\|f_{S(t)} - f_{S^*(t)}\|_2 \leq \lambda t e^{-\lambda t} \|\theta_X - \theta_Y\|_2$ , where  $\theta_Y(s) = \sum_{j=1}^n \omega_j \mu_j / (\mu_j + s \sqrt{-1})$ .

*Proof.* Using the Hausdorff-Young Theorem, one may conclude that  $\|f_{S(t)} - f_{S^*(t)}\|_2 \leq \|e^{\lambda t(\theta_X - 1)} - e^{\lambda t(\theta_Y - 1)}\|_2$ .

The rest of proof arrives by using the fact that  $\psi_X$  and  $\theta_Y$  are (or can be extend to) two meromorphic functions.  $\square$

### 3 Ruin Probability

This section utilizes integro-differential Equations (3) and (4) to derive an approximate formula for the infinite (and finite)-time ruin probability of a compound Poisson process (1). We seek an analytical solution  $\tilde{\psi}(\cdot)$  which is an exponential type function. In the other word, we assume:  $|\tilde{\psi}(\omega)| \leq M e^{T|\omega|}$ ,  $\omega \in \mathbb{C}$ , for some real numbers  $M$  and  $T$  in  $\mathbb{R}$ .

If this assumption is not met, as might be the case if, for example, there are point masses in  $\psi(\cdot)$ , our method works, but our error bounds may not be valid anymore.

The following theorem provides an  $(n + 1)$ -order ODE for infinite-time ruin probability  $\psi(\cdot)$  in the situation that claim size distribution  $X$  has been approximated by an  $n$ -component Hyperexponential density function  $f_X^*(\cdot)$ .

**Theorem 1.** Suppose claim size density function  $f_X(\cdot)$  has been approximated by an  $n$ -component Hyperexponential density function  $f_X^*(\cdot)$ . Then, infinite-time survival probability  $\tilde{\psi}(\cdot)$  of a compound Poisson process (1) can be approximated by infinite-time survival probability  $\tilde{\psi}_*(\cdot)$  which can be evaluated using the following  $(n + 1)$ -order ODE.  $\sum_{i=1}^n \lambda \omega_i \mu_i \tilde{\psi}_*^{(n-1)}(u) + \sum_{i=1}^n \sum_{i \neq j} \lambda \omega_i \mu_i \mu_j \tilde{\psi}_*^{(n-2)}(u) - \sum_{i=1}^n \sum_{i \neq j} \sum_{k > j, \neq i} \lambda \omega_i \mu_i \mu_j \mu_k \tilde{\psi}_*^{(n-3)}(u) + \sum_{i=1}^n \sum_{i \neq j} \sum_{k > j, \neq i} \sum_{l > k, \neq i} \lambda \omega_i \mu_i \mu_j \mu_k \mu_l \tilde{\psi}_*^{(n-4)}(u) + \dots + (-1)^n \sum_{i=1}^n \prod_{j \neq i} \mu_j \tilde{\psi}_*^{(0)}(u) - [\lambda \tilde{\psi}_*^{(n)}(u) - c \tilde{\psi}_*^{(n+1)}(u)] - \sum_{i=1}^n \mu_i [\lambda \tilde{\psi}_*^{(n-1)}(u) - c \tilde{\psi}_*^{(n)}(u)] - \sum_{i=1}^n \sum_{j \neq i} \mu_i \mu_j [\lambda \tilde{\psi}_*^{(n-2)}(u) - c \tilde{\psi}_*^{(n-1)}(u)] - \sum_{i=1}^n \sum_{j \neq i} \sum_{k > j, \neq i} \mu_i \mu_j \mu_k [\lambda \tilde{\psi}_*^{(n-3)}(u) - c \tilde{\psi}_*^{(n-2)}(u)] - \dots - \prod_{i=1}^n \mu_i [\lambda \tilde{\psi}_*^{(0)}(u) - c \tilde{\psi}_*^{(1)}(u)] = 0$  with boundary conditions that satisfy  $c \tilde{\psi}_*^{(m)}(0) - \lambda \tilde{\psi}_*^{(m-1)}(0) + \lambda \sum_{j=0}^{m-2} \tilde{\psi}_*^{(j)}(0) f^{(m-2-j)}(0) = 0$ , for  $m = 1, \dots, n$ .

*Proof.* An application of Lemma (2) by changing  $k(u) \mapsto -c \tilde{\psi}_*^{(1)}(u) + \lambda \tilde{\psi}_*(u)$ ,  $y(u) \mapsto \tilde{\psi}_*(u)$ , and  $\omega_i \mapsto \lambda \omega_i$  lead to the desired result.  $\square$

Using the fact that  $\tilde{\psi}_*(0) = 1 - \lambda E(X)/c$ , (see [10], Page

104) the above boundary condition equation leads to:

$$\begin{aligned} \tilde{\psi}_*^{(1)}(0) &= \tilde{\psi}_*(0) \frac{\lambda}{c}, \\ \tilde{\psi}_*^{(2)}(0) &= \tilde{\psi}_*(0) \left[ \left(\frac{\lambda}{c}\right)^2 - \left(\frac{\lambda}{c}\right) f_X(0) \right], \\ \tilde{\psi}_*^{(3)}(0) &= \tilde{\psi}_*(0) \left[ \left(\frac{\lambda}{c}\right)^3 - 2 f_X(0) \left(\frac{\lambda}{c}\right)^2 - \left(\frac{\lambda}{c}\right) f_X^{(1)}(0) \right], \end{aligned}$$

and so on.

The following provides error bound for approximating infinite-time survival probability  $\tilde{\psi}(\cdot)$  by  $\tilde{\psi}_*(\cdot)$ .

**Theorem 2.** Suppose claim size density function  $f_X(\cdot)$  has been approximated by an  $n$ -component Hyperexponential density function  $f_X^*(\cdot)$ . Then, the infinite-time survival probability  $\tilde{\psi}(u)$  of compound Poisson process (1) can be approximated by  $\tilde{\psi}_*(u)$ , given by Theorem (1), and its error satisfies  $\|\psi(u) - \psi_*(u)\|_2 \leq \frac{c \lambda \tilde{\psi}(0)}{\sqrt{\pi} a_1} \left\| \varphi_X(s) - \sum_{j=1}^n \frac{\omega_j \mu_j}{\mu_j + s} \right\|_2$ , where  $a_1 = \sup\{\varphi_X(s), \sum_{j=1}^n \frac{\omega_j \mu_j}{\mu_j + s}\}$  and  $\varphi_X(s)$  stands for the characteristic function of random claim  $X$ .

*Proof.* Application of the Hausdorff-Young for Laplace transform (Lemma 1) along with fact that  $\mathcal{L}(g'(x); x; s) = s \mathcal{L}(g(x); x; s) - g(0)$  and  $\mathcal{L}(\int_0^x (g(x-y)f(y)dy; x; s) = \mathcal{L}(g(x); x; s) \mathcal{L}(f(x); x; s)$ , one may conclude that

$$\begin{aligned} \|\psi(u) - \psi_*(u)\|_2 &\leq \frac{1}{\sqrt{\pi}} \|\mathcal{L}(\tilde{\psi}) - \mathcal{L}(\tilde{\psi}_*)\|_2 \\ &= \frac{1}{\sqrt{\pi}} \left\| \frac{c \tilde{\psi}(0)}{cu - \lambda + \lambda \mathcal{L}(f)} - \frac{c \tilde{\psi}(0)}{cu - \lambda + \lambda \mathcal{L}(f^*)} \right\|_2. \end{aligned}$$

Application of inequality  $\|1/h_1 - 1/h_2\|_2 \leq a^{-2} \|h_1 - h_2\|_2$ , where  $a = \sup\{h_1, h_2\}$ , from [11] completes the desired proof.  $\square$

The following theorem provides an  $(n + 1)$ -order PDE for finite-time ruin probability  $\psi(\cdot)$  in the situation that claim size distribution  $X$  has been approximated by an  $n$ -component Hyperexponential distribution function.

**Theorem 3.** Suppose claim size density function  $f_X(\cdot)$  has been approximated by an  $n$ -component Hyperexponential density function  $f_X^*(\cdot)$ . Then, finite-time survival probability  $\tilde{\psi}(u; T)$  of a compound Poisson process (1) can be approximated by finite-time survival probability  $\tilde{\psi}_*(u; T)$  which can be evaluated using the following

$(n + 1)$ -order PDE.

$$\begin{aligned}
 0 &= \sum_{i=1}^n \lambda \omega_i \mu_i \frac{\partial^{n-1}}{\partial u^{n-1}} \tilde{\psi}_*(u; T) + \sum_{i=1}^n \sum_{i \neq j}^n \lambda \omega_i \mu_i \mu_j \frac{\partial^{n-2}}{\partial u^{n-2}} \tilde{\psi}_*(u; T) \\
 &\quad - \sum_{i=1}^n \sum_{i \neq j}^n \sum_{k > j, \neq i}^n \lambda \omega_i \mu_i \mu_j \mu_k \frac{\partial^{n-3}}{\partial u^{n-3}} \tilde{\psi}_*(u; T) \\
 &\quad + \sum_{i=1}^n \sum_{i \neq j}^n \sum_{k > j, \neq i}^n \sum_{l > k, \neq i}^n \lambda \omega_i \mu_i \mu_j \mu_k \mu_l \frac{\partial^{n-4}}{\partial u^{n-4}} \tilde{\psi}_*(u; T) \\
 &\quad - \dots + (-1)^n \sum_{i=1}^n \prod_{j \neq i}^n \mu_j \frac{\partial^0}{\partial u^0} \tilde{\psi}_*(u; T) \\
 &\quad - \left[ \lambda \frac{\partial^n}{\partial u^n} \tilde{\psi}_*(u; T) - c \frac{\partial^{n+1}}{\partial u^{n+1}} \tilde{\psi}_*(u; T) + c \frac{\partial^{n+1}}{\partial T \partial u^n} \tilde{\psi}_*(u; T) \right] \\
 &\quad - \sum_{i=1}^n \mu_i \left[ \lambda \frac{\partial^{n-1}}{\partial u^{n-1}} \tilde{\psi}_*(u; T) - c \frac{\partial^n}{\partial u^n} \tilde{\psi}_*(u; T) + c \frac{\partial^n}{\partial T \partial u^{n-1}} \tilde{\psi}_*(u; T) \right] \\
 &\quad - \sum_{i=1}^n \sum_{j \neq i}^n \mu_i \mu_j \left[ \lambda \frac{\partial^{n-2}}{\partial u^{n-2}} \tilde{\psi}_*(u; T) - c \frac{\partial^{n-1}}{\partial u^{n-1}} \tilde{\psi}_*(u; T) \right. \\
 &\quad \quad \left. + c \frac{\partial^{n-1}}{\partial T \partial u^{n-2}} \tilde{\psi}_*(u; T) \right] \\
 &\quad - \sum_{i=1}^n \sum_{j \neq i}^n \sum_{k > j}^n \mu_i \mu_j \mu_k \left[ \lambda \frac{\partial^{n-3}}{\partial u^{n-3}} \tilde{\psi}_*(u; T) - c \frac{\partial^{n-2}}{\partial u^{n-2}} \tilde{\psi}_*(u; T) \right. \\
 &\quad \quad \left. + c \frac{\partial^{n-2}}{\partial T \partial u^{n-3}} \tilde{\psi}_*(u; T) \right] \\
 &\quad - \dots - \prod_{i=1}^n \mu_i \left[ \lambda \frac{\partial^0}{\partial u^0} \tilde{\psi}_*(u; T) - c \frac{\partial^1}{\partial u^1} \tilde{\psi}_*(u; T) + c \frac{\partial^1}{\partial T^1} \tilde{\psi}_*(u; T) \right],
 \end{aligned}$$

where  $\tilde{\psi}_*^{(n)}(0; T) = \lim_{u \rightarrow 0} \frac{\partial^n}{\partial u^n} \tilde{\psi}_*(u; T)$  and boundary conditions that satisfy  $c\tilde{\psi}_*^{(m)}(0; T) - c \frac{\partial}{\partial T} \tilde{\psi}_*^{(m-1)}(0; T) - \lambda \tilde{\psi}_*^{(m-1)}(0; T) + \lambda \sum_{j=0}^{m-2} \tilde{\psi}_*^{(j)}(0; T) f_X^{(n-2-j)}(0) = 0$ , for  $m = 1, \dots, n$ .

*Proof.* Using partial integro-differential equation (4) and the Schwarz integrability condition, one may change order of differentiation and obtain the above recursive formula for boundary conditions. An application of Lemma (2) by changing  $k(u) \mapsto -c \frac{\partial}{\partial u} \tilde{\psi}_*(u; T) + c \frac{\partial}{\partial T} \tilde{\psi}_*(u; T) + \lambda \tilde{\psi}_*(\cdot)$ ,  $y(u) \mapsto \tilde{\psi}_*(u; T)$ , and  $\omega_i \mapsto \lambda \omega_i$  lead to the desired result.  $\square$

Using the fact that  $\tilde{\psi}_*(u; 0) = 1$ ,  $\tilde{\psi}_*(0; T) = \int_0^{cT} F_{S,T}(x) dx / (cT)$ , and  $F_{S,T}(x) = P(\sum_{j=1}^{N(T)} X_j \leq x)$ , for all  $x \in \mathbb{R}^+$  (see [3], Page 121). One may compute the following from boundary conditions from recursive formula given by Theorem (3).

$$\begin{aligned}
 \tilde{\psi}_*^{(1)}(0; T) &= \frac{\lambda}{c} \tilde{\psi}_*(0; T) + \frac{\partial}{\partial T} \tilde{\psi}_*(0; T), \\
 \tilde{\psi}_*^{(2)}(0; T) &= \tilde{\psi}_*(0; T) \left[ \left(\frac{\lambda}{c}\right)^2 - \left(\frac{\lambda}{c}\right) f_X(0) \right] + \frac{\partial}{\partial T} \tilde{\psi}_*^{(1)}(0; T), \\
 \tilde{\psi}_*^{(3)}(0; T) &= \tilde{\psi}_*(0; T) \left[ \left(\frac{\lambda}{c}\right)^3 - 2f_X(0) \left(\frac{\lambda}{c}\right)^2 - \left(\frac{\lambda}{c}\right) f_X^{(1)}(0) \right] + \frac{\partial}{\partial T} \tilde{\psi}_*^{(2)}(0; T),
 \end{aligned}$$

where  $\tilde{\psi}_*^{(n)}(0; T) = \lim_{u \rightarrow 0} \frac{\partial^n}{\partial u^n} \tilde{\psi}_*(u; T)$ .

Using the central limit theorem for compound sum  $N(t)$   $\sum_{i=1}^n X_i$  (see [10], §2.5, or [27], §1.9), one may provide the following approximation for expression  $\tilde{\psi}_*(0; T) = \int_0^{cT} F_{S,T}(x) dx / (cT)$

$$\tilde{\psi}_*(0; T) \approx \frac{1}{cT} \int_0^{cT} \Phi \left( \frac{x - \lambda T m_1}{\sqrt{\lambda T m_2}} \right) dx,$$

where  $m_i = E(X^i)$ , for  $i = 1, 2$ , and  $\Phi(\cdot)$  stands for cumulative distribution function for standard normal distribution, see [10], §2.5, or [27], §1.9, for other parametric approximation approaches and [24] for a nonparametric approximation approach. For heavy tailed random claim size  $X$  that the ordinary central limit theorem does not work properly. One has to employ an appropriated version of the central limit theorem, see [25] and [5], among others, for more details.

The following provides error bound for approximating finite-time survivals probability  $\tilde{\psi}(u; T)$  by  $\tilde{\psi}_*(u; T)$ .

**Theorem 4.** Suppose claim size density function  $f_X(\cdot)$  has been approximated by an  $n$ -component Hyperexponential density function  $f_X^*(\cdot)$ . Then, the infinite-time survival probability  $\tilde{\psi}(u; T)$  of compound Poisson process (1) can be approximated by  $\tilde{\psi}_*(u; T)$ , given by Theorem (3), and its error satisfies

$$\begin{aligned}
 \|Error\|_2 &\leq \frac{\lambda}{\sqrt{\pi}} \left[ -\frac{c\tilde{\psi}(0; T)}{a_1^2} + \frac{a_2 T}{c} - c\tilde{\psi}(0; T) a_1^2 a_2^2 a_3 \right] \\
 &\quad \times \left\| \varphi_X(s) - \sum_{j=1}^n \frac{\omega_j \mu_j}{\mu_j + s} \right\|_2,
 \end{aligned}$$

where  $Error = \psi(u; T) - \psi_*(u; T)$ ,  $a_1 = \sup\{\varphi_X(s), \sum_{j=1}^n \frac{\omega_j \mu_j}{\mu_j + s}\}$ ,  $a_2 = \sup\{1/s - c/A(s), 1/s - c/A_*(s)\}$ ,  $a_3 = \sup\{e^{A(s)/cT}, e^{A_*(s)/cT}\}$ ,  $A(s) = cs - \lambda + \lambda \varphi_X(s)$ ,  $A_*(s) = cs - \lambda + \lambda \sum_{j=1}^n \omega_j \mu_j / (\mu_j + s)$ , and  $\varphi_X(s)$  stands for the characteristic function of random claim  $X$ .

*Proof.* Taking the Laplace transform from both sides of Equation (4) leads to the following first-order PDE

$$A(s) \mathcal{L}(\tilde{\psi}(u; T); u; s) - c\tilde{\psi}(0; T) - c \frac{\partial}{\partial T} \mathcal{L}(\tilde{\psi}(u; T); u; s) = 0,$$

where  $\mathcal{L}(\tilde{\psi}(u; 1); u; s) = 1/s$ . Therefore, the Laplace transform of finite-time ruin probability for compound Poisson process (1) is

$$\mathcal{L}(\tilde{\psi}(u; T); u; s) = \frac{c\tilde{\psi}(0; T)}{A(s)} + \left( \frac{1}{s} - \frac{c\tilde{\psi}(0; T)}{A(s)} \right) e^{A(s)/cT}.$$

The above finding along with an application of the Hausdorff-Young for Laplace transform (Lemma 1) lead to

$$\begin{aligned}
 \|E\|_2 &\leq \frac{1}{\sqrt{\pi}} \|\mathcal{L}(\tilde{\psi}(u; T); u; s) - \mathcal{L}(\tilde{\psi}_*(u; T); u; s)\|_2 \\
 &= \frac{1}{\sqrt{\pi}} \left\| \frac{c\tilde{\psi}(0; T)}{A(s)} + \left( \frac{1}{s} - \frac{c\tilde{\psi}(0; T)}{A(s)} \right) e^{A(s)/cT} \right. \\
 &\quad \left. - \frac{c\tilde{\psi}_*(0; T)}{A_*(s)} - \left( \frac{1}{s} - \frac{c\tilde{\psi}_*(0; T)}{A_*(s)} \right) e^{A_*(s)/cT} \right\|_2 \\
 &\leq \frac{c\tilde{\psi}(0; T)}{\sqrt{\pi} a_1^2} \|A(s) - A_*(s)\|_2 + \frac{b}{\sqrt{\pi}} \left\| \frac{TA(s)}{c} - \frac{TA_*(s)}{c} \right\|_2 \\
 &\quad + \frac{b}{\sqrt{\pi}} \left\| \ln \left( \frac{1}{s} - \frac{c\tilde{\psi}(0; T)}{A(s)} \right) - \ln \left( \frac{1}{s} - \frac{c\tilde{\psi}_*(0; T)}{A_*(s)} \right) \right\|_2,
 \end{aligned}$$

where  $E = \psi(u; T) - \psi_*(u; T)$ , the second inequality arrives by application of inequality  $\|1/h_1 - 1/h_2\|_2 \leq$

$a^{-2} \|h_1 - h_2\|_2$ , and  $a = \sup\{h_1, h_2\}$ , from [11], triangle inequality, and the Mean value theorem (i.e.,  $(\exp\{A(s)/cT + \ln(\frac{1}{s} - \frac{c\tilde{\psi}(0;T)}{A(s)})\} - \exp\{A_*(s)/cT + \ln(\frac{1}{s} - \frac{c\tilde{\psi}(0;T)}{A_*(s)})\}) / (A(s)/cT + \ln(\frac{1}{s} - \frac{c\tilde{\psi}(0;T)}{A(s)}) - A_*(s)/cT - \ln(\frac{1}{s} - \frac{c\tilde{\psi}(0;T)}{A_*(s)})) \leq b$  where  $b = \sup\{A(s)/cT + \ln(\frac{1}{s} - \frac{c\tilde{\psi}(0;T)}{A(s)}), A_*(s)/cT + \ln(\frac{1}{s} - \frac{c\tilde{\psi}(0;T)}{A_*(s)})\}$ ).

Application of inequality  $|\ln h_1 - \ln h_2| \leq \|h_1 - h_2\|_2 / a$ , where  $a = \sup\{h_1, h_2\}$ , from [11] completes the desired proof.  $\square$

### 4 Simulation Study

Consider compound Poisson process (1) with intensity rate  $\lambda = 1$  and premium  $c = 1.1$ . This section conducts two simulation studies to show practical application the about findings.

**Example 1.** Suppose random claim  $X$  in compound Poisson process (1) has been distributed according to Weibull(0.3,9.26053). Ref. [7] using a three-moment matching algorithm showed that density function of random claim  $X$  can be approximated by the following 2-component Hyperexponential density function

$$f_X^*(x) = 0.000095e^{-0.019x} + 1.348225e^{-1.355x}. \quad (7)$$

For infinite-time ruin probability: Application of Theorem (1) leads to the following second order ODE

$$1.1\tilde{\psi}_*^{(3)}(u) + 0.5114\tilde{\psi}_*^{(2)}(u) + 0.0026395\tilde{\psi}_*^{(1)}(u) = 0$$

with initial conditions  $\tilde{\psi}_*(0) = 0.0909090909$ ,  $\lim_{u \rightarrow 0} \tilde{\psi}_*^{(1)}(u) = 0.08264462809$ , and  $\lim_{u \rightarrow 0} \tilde{\psi}_*^{(2)}(u) = -0.03629992491$ .

Solving the above ODE, one may approximate finite-time survival probability  $\tilde{\psi}(u)$  of compound Poisson process (1) by  $\tilde{\psi}_*(u) = 0.9974815963 - 0.2101123939e^{-0.01502369720u} - 0.2873692025e^{-0.8589763028u}$ .

Figure 1 illustrates behavior for such approximated infinite-time ruin probability.

For finite-time ruin probability: Application of Theorem (3) leads to the following PDE  $1.1\frac{\partial^3}{\partial u^3}\tilde{\psi}_*(u;T) + 0.5114\frac{\partial^2}{\partial u^2}\tilde{\psi}_*(u;T) - 0.0539995\frac{\partial}{\partial u}\tilde{\psi}_*(u;T) - 1.1\frac{\partial^3}{\partial u^3}\tilde{\psi}_*(u;T) - 1.5114\frac{\partial^2}{\partial u^2}\tilde{\psi}_*(u;T) - 0.0283195\frac{\partial}{\partial T}\tilde{\psi}_*(u;T) = 0$  with initial conditions  $\tilde{\psi}_*(u,0) = 0$ ,  $\tilde{\psi}_*(0;T) = \beta(T)$ ,  $\lim_{u \rightarrow 20} \tilde{\psi}_*(u;T) = 1$ ,  $\lim_{u \rightarrow 0} \frac{\partial}{\partial u}\tilde{\psi}_*(u;T) = 0.9091\beta(T) + \frac{\partial}{\partial T}\beta(T)$ , where  $\beta(T) = \frac{1}{1.1T} \int_0^{1.1T} \Phi\left(\frac{x-T}{\sqrt{29.36T}}\right) dx$ .

Solving the above PDE, one may approximate finite-time survival probability  $\tilde{\psi}(u;T)$  of compound Poisson process (1) by  $\tilde{\psi}_*(u;T)$ , that its behavior (for  $T = 50, 100, 200$ ) has been illustrated in Figure 2.




Figure 1: Behavior for approximated: infinite-time ruin probability  $\psi_*(u)$ .




Figure 2: Behavior for approximated finite-time ruin probability  $\psi_*(u;T)$ , for  $T = 50, 100, 200$ .

**Example 2.** Suppose random claim  $X$  in compound Poisson process (1) has been distributed according to Gamma(0.7310,1). Ref. [29] using the Padé approximant method showed that density function of random claim  $X$  can be approximated by the following 3-component Hyperexponential density function

$$f_X^*(x) = 0.8099e^{-3.2398x} + 0.3616e^{-1.4465x} + 0.5198e^{-1.0396x}. \quad (8)$$

For infinite-time ruin probability: Application of Theorem (1) leads to the following second order ODE  $1.1\tilde{\psi}_*^{(4)}(u) + 5.29849\tilde{\psi}_*^{(3)}(u) + 6.479507012\tilde{\psi}_*^{(2)}(u) + 1.797919457\tilde{\psi}_*^{(1)}(u) = 0$  with initial conditions  $\tilde{\psi}_*(0) = 0.3354861821$ ,  $\lim_{u \rightarrow 0} \tilde{\psi}_*^{(1)}(u) = 0.3049874383$ ,  $\lim_{u \rightarrow 0} \tilde{\psi}_*^{(2)}(u) = -0.2385743202$ , and  $\lim_{u \rightarrow 20} \tilde{\psi}_*(u) = 1$ .

Solving the above ODE, one may approximate infinite-time survival probability  $\tilde{\psi}(u)$  of compound Poisson process (1) by  $\tilde{\psi}_*(u) = 1 - 0.013037e^{-3.073097u} - 0.008568e^{-1.349630u} - 0.642908e^{-0.394082u}$ . Figure 3 illustrates behavior for such approximated infinite-time ruin probability.

For finite-time ruin probability: Application of Theorem (3) leads to the following PDE  $1.1\frac{\partial^4}{\partial u^4}\tilde{\psi}_*(u;T) + 5.29849\frac{\partial^3}{\partial u^3}\tilde{\psi}_*(u;T) + 6.479507012\frac{\partial^2}{\partial u^2}\tilde{\psi}_*(u;T) + 1.797919457\frac{\partial}{\partial u}\tilde{\psi}_*(u;T) - 5.359146\frac{\partial}{\partial T}\tilde{\psi}_*(u;T) = 0$

$$10.5141 \frac{\partial^2}{\partial u \partial T} \tilde{\psi}_*(u; T) - 6.2985 \frac{\partial^3}{\partial u^2 \partial T} \tilde{\psi}_*(u; T) - 1.1 \frac{\partial^4}{\partial u^3 \partial T} \tilde{\psi}_*(u; T) = 0$$

with initial conditions  $\tilde{\psi}_*(u, 0) = 0$ ,  $\tilde{\psi}_*(0; T) = \beta(T)$ ,  $\lim_{u \rightarrow 20} \tilde{\psi}_*(u; T) = 1$ ,  $\lim_{u \rightarrow 0} \frac{\partial}{\partial u} \tilde{\psi}_*(u; T) = 0.9091\beta(T) + \frac{\partial}{\partial T}\beta(T)$ ,  $\lim_{u \rightarrow 0} \frac{\partial}{\partial u} \tilde{\psi}_*(u; T) = -0.71113\beta(T) + 0.9091 \frac{\partial}{\partial T}\beta(T) + \frac{\partial^2}{\partial T^2}\beta(T)$ , where  $\beta(T) = \frac{1}{1.1T} \int_0^{1.1T} \Phi\left(\frac{x-0.7309651999T}{\sqrt{0.7309651995T}}\right) dx$ .

Solving the above PDE, one may approximate finite-time survival probability  $\tilde{\psi}(u; T)$  of compound Poisson process (1) by  $\tilde{\psi}_*(u; T)$ , that its behavior (for  $T = 50, 100, 200$ ) has been illustrated in Figure 4.




Figure 3: Behavior for approximated infinite-time ruin probability  $\psi_*(u)$ .




Figure 4: Behavior for approximated finite-time ruin probability  $\psi_*(u; T)$ , for  $T = 50, 100, 200$ .

It worthwhile to mention that: A given density function (or a density function corresponding to a given data set) can be approximated by a Hyperexponential distribution using a Matlab package called “bayesf”, see [28] for more details.

## 5 Conclusion and Suggestions

This article approximates claim size density function  $f_X(\cdot)$  by a  $n$ -component Hyperexponential density function  $f_X^*(\cdot)$ . Then, it restates the problem of finding an infinite-time (or finite-time) ruin probability as a  $(n + 1)$ -order ordinary differential equation (or a partial differential equation for finite-time ruin probability).

Application of our findings has been given though a simulation study.

Certainly the following generalized Hyperexponential distribution can be provided a more accurate approximation in the situation that the true density function (or recorded data) has more than one mode.

$$g_X^{GHE}(x) = \sum_{i=1}^n \omega_i \mu_i e^{-\mu_i(x-b_i)} I_{[b_i, \infty)}(x). \quad (9)$$

In such situation the finite and infinite ruin probabilities can be evaluated using the following lemma.

**Lemma 4.** Suppose claim size density function  $f_X(\cdot)$  has been approximated by generalized Hyperexponential distribution  $g_X^{GHE}(\cdot)$ . The survival probability can be found by the following two inverse Laplace transforms.

- (i) The Laplace transform of the infinite-time survival probability can be found by the following equation

$$\mathcal{L}(\tilde{\psi}(u); u; s) = \frac{c\tilde{\psi}(0)}{cs - \lambda + \lambda \sum_{i=1}^k \frac{\omega_i \mu_i}{\mu_i + s} e^{-sb_i}}$$

- (ii) The Laplace transform of the finite-time survival probability can be found by the following equation

$$\mathcal{L}(\tilde{\psi}(u; T); u; s) = \frac{c\tilde{\psi}(0; T)}{B(s)} + \left(\frac{1}{s} - \frac{c\tilde{\psi}(0; T)}{B(s)}\right) e^{-\frac{B(s)}{cT}},$$

where  $B(s) = cs - \lambda + \lambda \sum_{i=1}^k \frac{\omega_i \mu_i}{\mu_i + s} e^{-sb_i}$ .

*Proof.* The desired result arrives by taking a Laplace transform from both sides of equations (3) and (4) and solving corresponding first-order PDE with boundary condition  $\tilde{\psi}(u; 0) = 1$  or  $\mathcal{L}(\tilde{\psi}(u; 0); u; s) = 1/s$ .  $\square$  Another possibility can be using a Lévy process to evaluate the ruin probability, see [21] and [22] for more details.

## Acknowledgements

Authors would like to thank professor Søren Asmussen for his constrictive comments on this article.

## References

- [1] Albrecher, H., Teugels, J. L., & Tichy, R. F. On a gamma series expansion for the time-dependent probability of collective ruin. *Insurance: Mathematics and Economics*, vol. 29, no. 3, pp. 345-355, 2001.
- [2] Avram, F., Chedom, D. F., & Horváth, A. On moments based Padé approximations of ruin probabilities. *Journal of computational and applied mathematics*, vol. 235, no. 10, pp. 3215-3228, 2011.
- [3] Asmussen, S., & Albrecher, H. *Ruin probabilities* (vol. 14). World Scientific, New Jersey, 2010.
- [4] Belding, D. F., & Mitchell, K. J. *Foundations of analysis* (vol. 79). Courier Corporation, New York, 2008.

- [5] Bilik, A. *Heavy-tail central limit theorem*. Lecture notes from University of California, San Diego, 2008.
- [6] Champeney, D. C. *A handbook of Fourier theorems*. Cambridge University press, New York, 1987.
- [7] Feldmann, A., & Whitt, W. Fitting mixtures of exponentials to long-tail distributions to analyze network performance models. *Performance evaluation*, vol. 31, no. 3, pp. 245-279, 1998.
- [8] Feller, W. *An introduction to probability and its applications*, vol. II. Wiley, New York, 1971.
- [9] Jewell, N. P. Mixtures of Exponential Distributions. *The Annals of Statistics*, vol. 10, no. 2, pp. 479-484, 1982.
- [10] Kaas, R., Goovaerts, M., Dhaene, J., & Denuit, M. *Modern actuarial risk theory: using R* (vol. 128). Springer-Verlag, Berlin Heidelberg, 2008.
- [11] Kucerovsky, D., & Payandeh Najafabadi, A. T. An approximation for a subclass of the Riemann-Hilbert problems. *IMA journal of applied mathematics*, vol. 74, no. 4, pp. 533-547, 2009.
- [12] Kucerovsky, D. Z. & Payandeh Najafabadi, A. T. Solving an Integral Equation Arising from the Ruin Probability of Long-term Bonus-Malus Systems. arXiv preprint arXiv:1701.05570. 2017.
- [13] Lukacs, E. *Characteristic Functions*. Oxford University Press, New York, 1987.
- [14] Makroglou, A. Integral equations and actuarial risk management: some models and numerics. *Mathematical Modelling and Analysis*, vol. 8, no. 2, pp. 143-154, 2003.
- [15] Melnikov, A. *Risk analysis in finance and insurance*. CRC Press, 2011.
- [16] Merzon, A., & Sadov, S. Hausdorff-Young type theorems for the Laplace transform restricted to a ray or to a curve in the complex plane. *arXiv preprint arXiv:1109.6085*. 2011.
- [17] Pandey, J. *The Hilbert transform of Schwartz distributions and application*. John Wiley & Sons, New York, 1996.
- [18] Payandeh Najafabadi, Amir T. Kucerovsky D. A weak approximated solution for a subclass of Wiener-Hopf integral equation. *IAENG International Journal of Applied Mathematics*, vol. 39, no. 1, pp. 247-252, 2009.
- [19] Payandeh Najafabadi, Amir T. Kucerovsky D. A weak approximation for the Wiener-Hopf factorization. *Cogent Mathematics*, vol. 2, no. 1, pp. 1074-1077, 2015.
- [20] Payandeh Najafabadi, Amir T. and Kucerovsky, D. On Approximating Ruin Probability of Double Stochastic Compound Poisson Processes. arXiv preprint arXiv:1701.05536, 2017a.
- [21] Payandeh Najafabadi, Amir T. and Kucerovsky, D. On the Distribution of Extrema for a Class of Lévy Processes. arXiv preprint arXiv:1701.05568, 2017b.
- [22] Payandeh Najafabadi, Amir T. and Kucerovsky D. Weak Approximation for the Extrema's Distributions of Lévy Processes. arXiv preprint arXiv:1701.05466, 2017c.
- [23] Pervozvansky, A. A. Equation for survival probability in a finite time interval in case of non-zero real interest force. *Insurance: Mathematics and Economics*, vol. 23, no. 3, pp. 287-295, 1998.
- [24] Qin, L., & Pitts, S. M. Nonparametric estimation of the finite-time survival probability with zero initial capital in the classical risk model. *Methodology and Computing in Applied Probability*, vol. 14, no. 4, pp. 919-936, 2012.
- [25] Rachev, S. T. (Ed.). *Handbook of Heavy Tailed Distributions in Finance: Handbooks in Finance* (vol. 1). Elsevier, Netherlands, 2003.
- [26] Sadov, S. Y. Characterization of Carleson measures by the Hausdorff-Young property. *Mathematical Notes*, vol. 94, no. 3, pp. 551-558, 2013.
- [27] Schmidli, H. *Lecture notes on Risk Theory*. Institute of Mathematics University of Cologne, German, 2006.
- [28] Sylvia, F. S. *Finite mixture and Markov switching models: Implementation in MATLAB using the package bayesf Version 2.0*. Springer-Verlag, New York, 2008.
- [29] Tran, D. X. Padé approximants for finite time ruin probabilities. *Journal of Computational and Applied Mathematics*, vol. 278, no. 1, pp. 130-137, 2015.
- [30] Walnut, D. F. *An introduction to wavelet analysis*. 2nd ed. Birkhäuser publisher, New York, 2002.