

Minimum Degree of Graphs and (g, f, n) -Critical Graphs

Sizhong Zhou *

Abstract—Let G be a graph of order p , and let a and b and n be nonnegative integers with $1 \leq a \leq b$, and let g and f be two integer-valued functions defined on $V(G)$ such that $a \leq g(x) \leq f(x) \leq b$ for all $x \in V(G)$. A (g, f) -factor of graph G is defined as a spanning subgraph F of G such that $g(x) \leq d_F(x) \leq f(x)$ for each $x \in V(G)$. Then a graph G is called a (g, f, n) -critical graph if after deleting any n vertices of G the remaining graph of G has a (g, f) -factor. In this paper, we prove that every graph G is a (g, f, n) -critical graph if its minimum degree is greater than $p+a+b-2\sqrt{(a+1)p-bn+1}$. Furthermore, it is showed that the result in this paper is best possible in some sense.

Keywords: graph, minimum degree, (g, f) -factor, (g, f, n) -critical graph

1 Introduction

In this paper, we consider a finite graph G with vertex set $V(G)$ and edge set $E(G)$, which has neither loops nor multiple edges. For any vertex x of G , we denote by $d_G(x)$ the degree of x in G . We denote by $\delta(G)$ the minimum vertex degree of G . For any $S \subseteq V(G)$, the subgraph of G induced by S is denoted by $G[S]$ and $G-S = G[V(G) \setminus S]$.

Let g and f be two nonnegative integer-valued functions defined on $V(G)$ such that $g(x) \leq f(x)$ for each $x \in V(G)$. A (g, f) -factor of graph G is defined as a spanning subgraph F of G such that $g(x) \leq d_F(x) \leq f(x)$ for each $x \in V(G)$ (Where of course d_F denotes the degree in F). If $g(x) = f(x)$ for each $x \in V(G)$, then a (g, f) -factor is called an f -factor. If $g(x) = a$ and $f(x) = b$ for all $x \in V(G)$, then a (g, f) -factor is called an $[a, b]$ -factor. If $g(x) = f(x) = k$ for all $x \in V(G)$, then a (g, f) -factor is called a k -factor. A graph G is called a (g, f, n) -critical graph if after deleting any n vertices of G the remaining graph of G has a (g, f) -factor. If G is a (g, f, n) -critical graph, then we also say that G is (g, f, n) -critical. If $g(x) = f(x)$ for each $x \in V(G)$, then a (g, f, n) -critical graph is an (f, n) -critical graph. If $g(x) = a$ and $f(x) =$

b for all $x \in V(G)$, then a (g, f, n) -critical graph is an (a, b, n) -critical graph. If $a = b = k$, then an (a, b, n) -critical graph is simply called a (k, n) -critical graph. In particular, a $(1, n)$ -critical graph is simply called an n -critical graph. The other notations and definitions not given in this paper can be found in [1].

Q. Yu [2] gave the characterization of n -critical graphs. O. Favaron [3] studied the properties of n -critical graphs. G. Liu and Q. Yu [4] studied the characterization of (k, n) -critical graphs. The characterization of (a, b, n) -critical graphs with $a < b$ was given by G. Liu and J. Wang [5]. S. Zhou [6-8] gave some sufficient conditions for graphs to be (a, b, n) -critical graphs. J. Li [9] showed two sufficient conditions for graphs to be (a, b, n) -critical graphs. S. Zhou [10] obtained a sufficient condition for graphs to be (g, f, n) -critical graphs. The characterization of (g, f, n) -critical graphs was given by J. Li and H. Matsuda [11]. In this paper, we obtain two new sufficient conditions for graphs to be (g, f, n) -critical graphs. The main results will be given in the following section. The following results on k -factors and (a, b, n) -critical graphs and (g, f, n) -critical graphs are known.

In [12], Y. Egawa and H. Enomoto proved the following result for the existence of k -factors.

Theorem 1 [12] *Let $k \geq 2$ be an integer, and let G be a graph of order n , kn is even. If*

$$\delta(G) > n + 2k - 2\sqrt{kn + 1},$$

then G has a k -factor.

In [5], G. Liu and J. Wang gave a necessary and sufficient condition for graphs to be (a, b, n) -critical graphs.

Theorem 2 [5] *Let a, b and n be nonnegative integers with $1 \leq a < b$, and G a graph of order $p \geq a + n + 1$. Then G is (a, b, n) -critical if and only if*

$$b|S| + d_{G-S}(T) - a|T| \geq bn, \text{ or} \\ \sum_{j=0}^{a-1} (a-j)p_j(G-S) \leq b|S| - bn$$

for all subset S of $V(G)$ with $|S| \geq n$, where $T = \{x :$

*Foundation items: This research was supported by Jiangsu Provincial Educational Department (07KJD110048). Received date: December 08, 2007. My address: School of Mathematics and Physics, Jiangsu University of Science and Technology, Mengxi Road 2, Zhenjiang, Jiangsu 212003, People's Republic of China. Email: zsz_cumt@163.com

$x \in V(G) \setminus S, d_{G-S}(x) \leq a - 1$ and $p_j(G - S) = |\{x \in V(G) \setminus S : d_{G-S}(x) = j\}|$.

In [10], S. Zhou showed a sufficient condition for graphs to be (g, f, n) -critical graphs.

Theorem 3 ^[10] *Let G be a graph, and let g and f be two nonnegative integer-valued functions defined on $V(G)$ such that $g(x) < f(x)$ for each $x \in V(G)$. If $g(x) \leq d_G(x)$ and $f(x)(d_G(y) - n) \geq d_G(x)g(y)$ for each $x, y \in V(G)$, then G is a (g, f, n) -critical graph. Here n is a nonnegative integer.*

2 The Proof of Main Theorems

In this paper, we obtain a new sufficient condition for graphs to be (g, f, n) -critical graphs. Our result is the extension of Theorem 1.

Theorem 4 *Let G be a graph of order p , and let a, b and n be nonnegative integers such that $1 \leq a < b$, and let g and f be two integer-valued functions defined on $V(G)$ such that $a \leq g(x) < f(x) \leq b$ for each $x \in V(G)$. If*

$$\delta(G) > p + a + b - 2\sqrt{(a+1)p - bn + 1}, \quad (1)$$

then G is a (g, f, n) -critical graph.

In Theorem 4, if $n = 0$, then we obtain the following corollary.

Corollary 1 *Let G be a graph of order p , and let a and b be integers such that $1 \leq a < b$, and let g and f be two integer-valued functions defined on $V(G)$ such that $a \leq g(x) < f(x) \leq b$ for each $x \in V(G)$. If*

$$\delta(G) > p + a + b - 2\sqrt{(a+1)p + 1},$$

then G has a (g, f) -factor.

According to Corollary 1 and the definition of (g, f, n) -critical graph, we obtain easily the following result.

Theorem 5 *Let G be a graph of order p , and let a, b and n be nonnegative integers such that $1 \leq a < b$, and let g and f be two integer-valued functions defined on $V(G)$ such that $a \leq g(x) < f(x) \leq b$ for each $x \in V(G)$. If*

$$\delta(G) > p + a + b - 2\sqrt{(a+1)p + 1} + n,$$

then G is a (g, f, n) -critical graph.

Let S and T be disjoint subsets of $V(G)$. We write $e_G(S, T) = |\{xy \in E(G) : x \in S, y \in T\}|$, $f(S) = \sum_{x \in S} f(x)$, $d_{G-S}(T) = \sum_{x \in T} d_{G-S}(x)$, and $g(T) = \sum_{x \in T} g(x)$. Our proof of Theorem 4 relies heavily on the following theorem.

Theorem 6 ^[11] *Let G be a graph, $n \geq 0$ an integer, and let g and f be two integer-valued functions defined on $V(G)$ such that $g(x) < f(x)$ for each $x \in V(G)$. Then G is a (g, f, n) -critical graph if and only if*

$$\begin{aligned} \delta_G(S, T) &= f(S) + d_{G-S}(T) - g(T) \\ &\geq \max\{f(N) : N \subseteq S, |N| = n\} \end{aligned}$$

for all disjoint subsets S and T of $V(G)$ with $|S| \geq n$.

Proof of Theorem 4. Suppose a graph G satisfies the condition of the theorem, but it is not a (g, f, n) -critical graph. Then, by Theorem 6, there exist disjoint subsets S and T of $V(G)$ with $|S| \geq n$ such that

$$\begin{aligned} \delta_G(S, T) &= f(S) + d_{G-S}(T) - g(T) \\ &\leq \max\{f(N) : N \subseteq S, |N| = n\} - 1. \end{aligned} \quad (2)$$

We choose subsets S and T such that $|T|$ is minimum and S and T satisfy (2).

Claim 1. $d_{G-S}(x) \leq g(x) - 1 \leq b - 2$ for each $x \in T$.

Proof. Suppose that there exists a vertex $x \in T$ such that $d_{G-S}(x) \geq g(x)$. Then the subsets S and $T - \{x\}$ satisfy (2), which contradicts the choice of T . Completing the proof of Claim 1.

If $T = \emptyset$, then by (2), $f(S) - 1 \geq \max\{f(N) : N \subseteq S, |N| = n\} - 1 \geq \delta_G(S, T) = f(S)$, a contradiction. Hence, $T \neq \emptyset$. Let

$$h = \min\{d_{G-S}(x) : x \in T\}.$$

According to Claim 1, we have

$$0 \leq h \leq b - 2,$$

and

$$\delta(G) \leq h + |S|. \quad (3)$$

According to (2) and $|S| + |T| \leq p$, we get that

$$\begin{aligned} bn - 1 &\geq \max\{f(N) : N \subseteq S, |N| = n\} - 1 \\ &\geq \delta_G(S, T) = f(S) + d_{G-S}(T) - g(T) \\ &\geq (a+1)|S| + d_{G-S}(T) - (b-1)|T| \\ &\geq (a+1)|S| + h|T| - (b-1)|T| \\ &= (a+1)|S| - (b-h-1)|T| \\ &\geq (a+1)|S| - (b-h-1)(p-|S|) \\ &= (a+b-h)|S| - (b-h-1)p. \end{aligned}$$

Thus, we obtain

$$|S| \leq \frac{(b-h-1)p + bn - 1}{a+b-h}. \quad (4)$$

In view of (3) and (4), we have

$$\delta(G) \leq h + |S| \leq h + \frac{(b-h-1)p + bn - 1}{a+b-h}. \quad (5)$$

Let $f(h) = h + \frac{(b-h-1)p+bn-1}{a+b-h}$. In the range of $h \leq a+b$, the function $f(h)$ attains its maximum value at $h = a+b - \sqrt{(a+1)p - bn + 1}$. Since $0 \leq h \leq b-2$, then we have

$$\begin{aligned} f(h) &\leq f(a+b - \sqrt{(a+1)p - bn + 1}) \\ &= p+a+b - 2\sqrt{(a+1)p - bn + 1}, \end{aligned}$$

that is,

$$\delta(G) \leq p+a+b - 2\sqrt{(a+1)p - bn + 1},$$

this contradicts (1).

From the argument above, we deduce the contradiction. Hence, G is a (g, f, n) -critical graph. Completing the proof of Theorem 4.

Remark. Let us show that the condition $\delta(G) > p+a+b - 2\sqrt{(a+1)p - bn + 1}$ in Theorem 4 can not be replaced by $\delta(G) \geq p+a+b - 2\sqrt{(a+1)p - bn + 1}$. Let $a = 2, b = 3$, and $n \geq 0$ an integer. Let $H = K_{n+1} \vee (K_a \cup K_a)$. Then $p = 2a + n + 1$ and $\delta(H) = a + n$. Thus, we obtain easily $p+a+b - 2\sqrt{(a+1)p - bn + 1} = a+n$, that is, $\delta(H) = p+a+b - 2\sqrt{(a+1)p - bn + 1}$. Let $S = V(K_{n+1}) \subseteq V(H)$, $T = V(K_a \cup K_a) \subseteq V(H)$. Since $a \leq g(x) < f(x) \leq b$ and $b = a+1$, then we have $g(x) = a$ and $f(x) = b = a+1$. Thus, we get

$$\begin{aligned} \delta_H(S, T) &= f(S) + d_{H-S}(T) - g(T) \\ &= b|S| + (a-1)|T| - a|T| \\ &= b|S| - |T| \\ &= b(n+1) - 2a \\ &= bn + b - 2a \\ &= bn - 1 \quad (\text{Since } a = 2 \text{ and } b = 3) \\ &< bn = \max\{f(N) : N \subseteq S, |N| = n\}. \end{aligned}$$

By Theorem 6, H is not a (g, f, n) -critical graph. In the above sense, the result of Theorem 4 is best possible.

We may adopt the similar way to argue the condition $\delta(G) > p+a+b - 2\sqrt{(a+1)p + 1 + n}$ in Theorem 5, and the condition $\delta(G) > p+a+b - 2\sqrt{(a+1)p + 1 + n}$ in Theorem 5 is the best possible in some sense.

References

- [1] J. A. Bondy, U. S. R. Murty, *Graph Theory with Applications*, London, The Macmillan Press, 1976.
- [2] Qinglin Yu, "Characterizations of various matching extensions in graphs," *Australasian Journal of Combinatorics*, V7, N5, pp. 55-64, 1/93
- [3] O. Favaron, "On k -factor-critical graphs," *Discussiones Mathematicae Graph Theory*, V16, N4, pp. 41-51, 1/96
- [4] Guizhen Liu, Qinglin Yu, " k -factors and extendability with prescribed components," *Congr. Numer.*, V139, N9, pp. 77-88, 1/99
- [5] Guizhen Liu, Jianfang Wang, " (a, b, k) -critical graphs," *Advances in Mathematics(China)*, V27, N6, pp. 536-540, 6/98
- [6] Sizhong Zhou, "Sufficient conditions for (a, b, k) -critical graphs," *Journal of Jilin University (Science Edition)(China)*, V43, N11, pp. 607-609, 5/05
- [7] Sizhong Zhou, Jiashang Jiang, "Notes on the binding numbers for (a, b, k) -critical graphs," *Bulletin of the Australian Mathematical Society*, V76, N5, pp. 307-314, 2/07
- [8] Sizhong Zhou, Minggang Zong, "Some new sufficient conditions for graphs to be (a, b, k) -critical graphs," *Ars Combinatoria*, to appear.
- [9] Jianxiang Li, "Sufficient conditions for graphs to be (a, b, n) -critical graphs," *Mathematica Applicata (China)*, V17, N22, pp. 450-455, 3/04
- [10] Sizhong Zhou, "Some sufficient conditions for graphs to have (g, f) -factors," *Bulletin of the Australian Mathematical Society*, V75, N8, pp. 447-452, 3/07
- [11] Jianxiang Li, H. Matsuda, "On (g, f, n) -critical graphs," *Ars Combinatoria*, V78, N7, pp. 71-82, 1/06
- [12] Y. Egawa, H. Enomoto, "Sufficient conditions for the existence of k -factors," *Recent Studies in Graph Theory*, Vishwa International Publication, India, pp. 96-105, 1/89