# Existence and Stability of Periodic Solution in Impulsive Hopfield Networks with Time-Varying Delays

Yanxia Cheng, Yan Yan, and Zhanji Gui\* Member, IAENG,

Abstract—By constructing suitable Lyapunov functions, we study the existence, uniqueness and global exponential stability of periodic solution for impulsive Hopfield neural networks with time-varying delays. Our condition extends and generalizes a known condition for the global exponential periodicity of continuous Hopfield neural networks with time-varying delays. Further the numerical simulation shows that our system can occur many forms of complexities including gui strange attractor and periodic solution.

*Index Terms*—Hopfield neural network, pulse, time-varying delay, Periodic solution, Numerical simulation.

#### I. INTRODUCTION

**T**N recent years, stability of different classes of neural networks with time delay, such as Hopfield neural networks, cellular neural networks, bidirectional associative neural networks, Lotka-Volterra neural networks, has been extensively studied and various stability conditions have been obtained for these models of neural networks [1]–[15].

Stability and convergence properties are generally regarded as important effects of delays. Both in biological and man-made neural systems, integration and communication delays are ubiquitous, and often become sources of instability. The delays in electronic neural networks are usually time varying, and sometimes vary violently with time due to the finite switching speed of amplifiers and faults in the electrical circuit. They slow down the transmission rate and tend to introduce some degree of instability in circuits. Therefore, fast response must be required in practical electronic neuralnetwork designs. The technique to achieve fast response troubles many circuit designers. So, it is important to investigate the delay independent stability and decay estimates of the states of analog neural networks.

However, besides delay effect, impulsive effect likewise exists in a wide variety of evolutionary processes in which states are changed abruptly at certain moments of time, involving such fields as medicine and biology, economics, mechanics, electronics and telecommunications, etc. Many interesting results on impulsive effect have been gained [16]– [30]. As artificial electronic systems, neural networks such as Hopfield neural networks, bidirectional neural networks and recurrent neural networks often are subject to impulsive

Y Cheng is with with the School of science, Beijing Forestry University, 100083, P.R. China.

Y. Yan and Z. Gui are with the School of Mathematics and Statistics, Hainan Normal University, Haikou, Hainan, 571158, P.R. China

\* Corresponding author e-mail:zhanjigui@sohu.com(Z. Gui)

perturbations which can affect dynamical behaviors of the systems just as time delays. Therefore, it is necessary to consider both impulsive effect and delay effect on the stability of neural networks.

In this paper, we consider the following impulsive Hopfield neural networks with time-varying delays:

$$\begin{cases} \dot{x}_{i}(t) = -a_{i}x_{i}(t) + \sum_{j=1}^{n} a_{ij}g_{j}(x_{j}(t)) \\ + \sum_{j=1}^{n} b_{ij}g_{j}(x_{j}(t - \tau_{ij}(t))) + I_{i}(t), \ t \neq t_{k}, \\ \Delta x_{i}(t_{k}) = \gamma_{ik}x_{i}(t_{k}), \ i = 1, 2, \dots, n, \ k = 1, 2, \dots, \end{cases}$$
(1)

where *n* is the number of neurons in the network,  $x_i(t)$  is the state of the *i*th neuron at time *t*,  $a_{ij}$  is the rate at which the *i*th neuron resets the state when isolated from the system,  $b_{ij}$  is the connection strength from the *j*th neuron to the *i*th neuron.  $g(x) = (g_1(x_1), g_2(x_2), \ldots, g_n(x_n))^T : \mathbb{R}^n \to \mathbb{R}^n$  is the output of the *i*th neuron at time *t*,  $I(t) = (I_1(t), I_2(t), \ldots, I_n(t))^T \in \mathbb{R}^n$  is the  $\omega$ -periodic external input to the *i*th neuron.

Throughout this paper, we assume that

(H<sub>1</sub>) For  $j \in \{1, ..., n\}$ ,  $g_j(u)$  (j = 1, 2, ..., n) is globally Lipschitz-continuous with the Lipschitz constant  $L_j > 0$ . That is,

$$|g_j(u_1) - g_j(u_2)| \leq L_j |u_1 - u_2|,$$

for all  $u_1, u_2 \in \mathbb{R} = (-\infty, \infty)$ .

- (H<sub>2</sub>) There exists a positive integer p such that,  $t_{k+p} = t_k + \omega, \gamma_{i(k+p)} = \gamma_{ik}, k > 0, k = 1, 2, \dots$
- (H<sub>3</sub>)  $\tau_{ij}(t)(i, j = 1, 2, ..., n)$  are continuously differentiable  $\omega$ -periodic functions defined on  $\mathbb{R}^+$ ,  $\tau = \sup_{0 \le t \le \omega} \tau_{ij}(t)$  and  $\inf_{t \in \mathbb{R}^+} \{1 - \dot{\tau}_i(t)\} > 0$ .

In order to describe the initial condition accompanying Eq. (1), we introduce the following notations.

Definition 1: A function  $\phi : [-\tau, 0] \to \mathbb{R}$  is said to be a  $C^*$ -function if the following two conditions are satisfied:

- (a)  $\phi$  is piecewise continuous with first kind discontinuity at the points  $t_k$ . Moreover,  $\phi$  is left-continuous at each discontinuity point.
- (b) For all  $i \in \{1, ..., n\}$  and  $k \in \{1, ..., p\}, \phi_i(t_k+0) = \phi_i(t_k) + \gamma_{ik}\phi_i(t_k)$ .

Let  $C^*$  denote the set of all the  $C^*$ -functions. Obviously,  $(C^*, \mathbb{R}, +, \cdot)$  forms a vector space on  $\mathbb{R}$ . Now consider  $(C^*, \mathbb{R}, +, \cdot)$  endowed with the norm defined by

$$\|\phi\|_{\infty} = \sup_{-\tau \leqslant \theta \leqslant 0} \|\phi(\theta)\| = \sup_{-\tau \leqslant \theta \leqslant 0} \max_{1 \leqslant i \leqslant n} |\phi_i(\theta)|.$$

Manuscript received March 6, 2012. This work was supported in part by the National Natural Science Foundation of China (60963025), Natural Sciences Foundation of Hainan Province (110007) and the Foundation of Hainan College of Software Technology (Hr200808, HrZD201101).

Proceedings of the World Congress on Engineering 2012 Vol I WCE 2012, July 4 - 6, 2012, London, U.K.

For brevity, let us denote this normed space by  $C^*$ . In this paper it is assumed that Eq. (1) is accompanied with the initial condition  $\phi \in C^*$ . We can easily prove.

Lemma 1:  $C^*$  is a Banach space.

Definition 2: A function  $x : [-\tau, \infty] \to \mathbb{R}$  is said to be the special solution of Eq. (1) with initial condition  $\phi \in C^*$ if the following two conditions are satisfied:

- (a) x is piecewise continuous with first kind discontinuity at the points  $t_k, k \in \{1, ..., p\}$ .
- (b) x satisfies Eq. (1) for  $t \ge 0$ , and  $x(\theta) = \phi(\theta)$  for  $\theta \in [-\tau, 0]$ .

Henceforth, we let  $x(t, \phi)$  denote the special solution of Eq. (1) with initial condition  $\phi \in C^*$ 

Definition 3: Eq. (1) is said to be globally exponentially periodic if (a) it possesses a periodic solution  $x(t, \phi^*)$ , and (b)  $x(t, \phi^*)$  is globally exponentially stable. That is, there exist positive constants  $\varepsilon$  and M such that every solution of Eq. (1) satisfies

$$\|x(t,\phi) - x(t,\phi^*)\|_{\infty} \leqslant M \|\phi - \phi^*\| e^{-\varepsilon t}, \text{ for all } t \ge 0.$$

In the next section, we will use the upper right Dini derivative of a continuous function f(t), which is defined by

$$D^+f(t) = \lim_{h \to 0^+} \sup_{0 < \Delta t \leqslant h} \frac{f(t + \Delta t) - f(t)}{\Delta t}.$$

The following lemma follows directly from the definition of upper right Dini derivative.

Lemma 2: Let f(t) be a continuous function on  $\mathbb{R}$  that is differentiable at  $t_0$ . Then

$$D^{+}|f(t_{0})| = \begin{cases} \dot{f}(t_{0}), & \text{when } f(t_{0}) > 0, \\ -\dot{f}(t_{0}), & \text{when } f(t_{0}) < 0, \\ |\dot{f}(t_{0})|, & \text{when } f(t_{0}) = 0. \end{cases}$$

### II. MAIN RESULT

Now we define  $\psi(t) = t - \tau_i(t)$ , then  $\psi^{-1}(t)$  has inverse function  $\nu$ . Set

$$\delta_i = \max\left\{\frac{1}{1 - \dot{\tau}_i(\psi_i^{-1}(t))} : t \in \mathbb{R}\right\}, i = 1, 2, \dots, n.$$

*Theorem 1:* Eq. (1) is globally exponentially periodic if the following two conditions are satisfied:

- (H<sub>4</sub>)  $|1 + \gamma_{ik}| \leq 1$ , for all  $i \in \{1, ..., n\}$ , and  $k \in \{1, ..., p\}$ ,
- (H<sub>5</sub>) There exist positive numbers  $\alpha_1, \alpha_2, \ldots, \alpha_n$ , such that

$$\alpha_i a_i > L_i \sum_{j=1}^n \alpha_j (|b_{ji}| + \delta_j |c_{ji}|), \quad i = 1, 2..., n.$$

As a direct result of Theorem 1, we have

Corollary 1: Eq. (1) is globally exponentially periodic if

(a)  $|1 + \gamma_{ik}| \leq 1$ , for all  $i \in \{1, ..., n\}$ , and  $k \in \{1, ..., p\}$ ,

(b)

$$a_i > L_i \sum_{j=1}^n (|b_{ji}| + \delta_j |c_{ji}|), \ i = 1, 2, \dots, n.$$

In order to prove Theorem 1, we need the following Lemma.

# ISBN: 978-988-19251-3-8 ISSN: 2078-0958 (Print); ISSN: 2078-0966 (Online)

Lemma 3: Let  $x(t, \phi)$ ,  $x(t, \varphi)$  be a pair of solutions of Eq. (1). If the two conditions given in Theorem 1 are satisfied, then there is a positive number  $\varepsilon$  such that,

$$\|x(t,\phi) - x(t,\varphi)\|_{\infty} \leqslant M(\varepsilon) \|\phi - \varphi\|_{\infty} e^{-\varepsilon t}, \text{ for all } t \ge 0$$
 where

$$M(\varepsilon) = \frac{1}{\min_{1 \le j \le n} \alpha_j} \sum_{i=1}^n \alpha_i \bigg[ 1 + \frac{1}{\varepsilon} L_j \delta_j |c_{ij}| (e^{\varepsilon \tau} - 1) \bigg].$$

**Proof.** Let  $x(t, \phi) = (x_1(t, \phi), x_2(t, \phi), \dots, x_n(t, \phi))^T$  and  $x(t, \varphi) = (x_1(t, \varphi), x_2(t, \varphi), \dots, x_n(t, \varphi))^T$  be an arbitrary pair of solutions of Eq. (1). Let

$$\Delta x_i(t,\phi,\varphi) = x_i(t,\phi) - x_i(t,\varphi)$$
  

$$\Delta g_j(x_j(t,\phi,\varphi)) = g_j(x_j(t,\phi)) - g_j(x_j(t,\varphi))$$
  

$$V(t) = \sum_{i=1}^n \alpha_i \bigg\{ |\Delta x_i(t,\phi,\varphi)| e^{-\varepsilon t}$$
  

$$+ \sum_{j=1}^n \int_{t-\tau_j(t)}^t \frac{L_j |c_{ij}|}{1 - \dot{\tau_j}(\psi_j^{-1}(s))}$$
  

$$\cdot |\Delta x_j(s,\phi,\varphi)| e^{\varepsilon(s+\tau_j(\psi_j^{-1}(s))} ds \bigg\}.$$
 (2)

We proceed by considering two possibilities.

**Case 1.**  $t \neq t_k$  for all  $k \in \{1, \ldots, p\}$ . From the second condition in Theorem 3, there is a small positive number  $\varepsilon$  such that

$$\alpha_i(a_i - \varepsilon) > L_i \sum_{j=1}^n \alpha_j \left( |b_{ji}| + \delta_j |c_{ji}| e^{\varepsilon \tau} \right), \tag{3}$$

where i = 1, ..., n. Calculating the derivatives of V(t) along the solutions of Eq. (1), we get

$$D^{+}V(t) = \sum_{i=1}^{n} \alpha_{i} \left\{ e^{-\varepsilon t} D^{+} |\Delta x_{i}(t,\phi,\varphi)| + \sum_{j=1}^{n} \left[ \frac{L_{j}|c_{ij}|}{1 - \dot{\tau}_{j}(\psi_{j}^{-1}(t))} |\Delta x_{j}(t,\phi,\varphi)| + e^{\varepsilon(t + \tau_{j}(\psi_{j}^{-1}(t)))} - L_{j}|c_{ij}| |\Delta x_{j}(t - \tau_{j}(t),\phi,\varphi)| e^{\varepsilon t} \right] + \varepsilon e^{-\varepsilon t} |\Delta x_{i}(t,\phi,\varphi)| \right\}.$$
(4)

Note that for  $i = 1, \ldots, n$ ,

$$\dot{x}_i(t,\phi) - \dot{x}_i(t,\varphi) = -a_i \Delta x_i(t,\phi,\varphi) + \sum_{j=1}^n b_{ij} \Delta g_j(x_j(t,\phi,\varphi)) + \sum_{j=1}^n c_{ij} \Delta g_j(x_j(t-\tau_j(t),\phi,\varphi)),$$

which plus  $(H_1)$  yields

D

$$|x(t,\phi) - x(t,\varphi)| \leq -a_i |\Delta x_i(t,\phi,\varphi)|$$

$$+ \sum_{j=1}^n b_{ij} |\Delta g_j(x_j(t,\phi,\varphi))|$$

$$+ \sum_{j=1}^n c_{ij} |\Delta g_j(x_j(t-\tau_j(t),\phi,\varphi))|$$

Proceedings of the World Congress on Engineering 2012 Vol I WCE 2012, July 4 - 6, 2012, London, U.K.

$$\leq -a_i |\Delta x_i(t,\phi,\varphi)| + \sum_{j=1}^n L_j |b_{ij}| |\Delta x_j(t,\phi,\varphi)|$$
  
+ 
$$\sum_{j=1}^n L_j |c_{ij}| |\Delta x_j(t-\tau_j(t),\phi,\varphi)|.$$
(5)

Substituting Eq. (5) into Eq. (4), we obtain

 $D^{\cdot}$ 

$${}^{+}V(t) = \sum_{i=1}^{n} \alpha_{i} \left[ -a_{i}e^{-\varepsilon t} |\Delta x_{i}(t,\phi,\varphi)| \right. \\ \left. +e^{-\varepsilon t} \sum_{j=1}^{n} L_{j} |b_{ij}| |\Delta x_{j}(t,\phi,\varphi)| \right. \\ \left. +e^{-\varepsilon t} \sum_{j=1}^{n} L_{j} |c_{ij}| |\Delta x_{j}(t-\tau_{j}(t),\phi,\varphi)| \right. \\ \left. +\varepsilon e^{-\varepsilon t} |\Delta x_{i}(t,\phi,\varphi)| \right. \\ \left. +\sum_{j=1}^{n} \frac{L_{j} |c_{ij}|}{1-\dot{\tau_{j}}(\psi_{j}^{-1}(t))} |\Delta x_{j}(t,\phi,\varphi)| \right. \\ \left. e^{\varepsilon (t+\tau_{j}(\psi_{j}^{-1}(t)))} -\sum_{j=1}^{n} L_{j} |c_{ij}| |\Delta x_{j}(t-\tau_{j}(t),\phi,\varphi)| e^{\varepsilon t} \right] \\ \leqslant e^{-\varepsilon t} \sum_{i=1}^{n} \alpha_{i} \left[ (\varepsilon - a_{i}) |\Delta x_{i}(t,\phi,\varphi)| \right. \\ \left. +\sum_{j=1}^{n} L_{j} |b_{ij}| |\Delta x_{j}(t,\phi,\varphi)| \right. \\ \left. +\sum_{j=1}^{n} L_{j} \delta_{j} |c_{ij}| e^{\varepsilon \tau} |\Delta x_{j}(t,\phi,\varphi)| \right] \\ \left. = e^{-\varepsilon t} \left[ \sum_{i=1}^{n} \alpha_{i}(\varepsilon - a_{i}) |\Delta x_{i}(t,\phi,\varphi)| \right. \\ \left. +\sum_{i=1}^{n} L_{i} \sum_{j=1}^{n} \alpha_{j} (|b_{ji}| + \delta_{i}|c_{ji}| e^{\varepsilon \tau}) \right] \right. \\ \left. \cdot |\Delta x_{i}(t,\phi,\varphi)| \\ \left. = e^{-\varepsilon t} \sum_{i=1}^{n} \left[ \alpha_{i}(\varepsilon - a_{i}) \right. \\ \left. +L_{i} \sum_{j=1}^{n} (|b_{ji}| + \delta_{i}|c_{ji}| e^{\varepsilon \tau}) \right] \right] \\ \left. \cdot |\Delta x_{i}(t,\phi,\varphi)| \leqslant 0.$$

**Case 2.**  $t = t_k$ , for some  $k \in \{1, 2, ..., p\}$ . Then

$$\begin{split} V(t+0) &= \sum_{i=1}^{n} \alpha_i \bigg[ |\Delta x_i(t+0,\phi,\varphi)| e^{-\varepsilon t} \\ &+ \sum_{j=1}^{n} \int_{t-\tau_j(t)}^{t} \frac{L_j |c_{ij}|}{1 - \dot{\tau_j}(\psi_j^{-1}(s))} \\ &\cdot \Delta x_j(s,\phi,\varphi) e^{\varepsilon(s+\tau_j(\psi_j^{-1}(s)))} ds \bigg]. \end{split}$$

According to Eq. (2) and  $(H_4)$ , we obtain

$$V(t+0) - V(t)$$
  
=  $e^{-\varepsilon t} \sum_{i=1}^{n} \alpha_i \Big( |\Delta x_i(t+0,\phi,\varphi)| - |\Delta x_i(t,\phi,\varphi)| \Big)$ 

$$= -e^{-\varepsilon t} \sum_{i=1}^{n} \alpha_i (1 - |1 + \gamma_{ik}|) |\Delta x_i(t, \phi, \varphi)| \leq 0.$$

Namely,  $V(t+0) \leq V(t)$ .

Combining the above discussions, we obtain  $V(t) \leq V(0)$ for all  $t \ge 0$ . This plus the inspections that

$$V(t) \ge e^{\varepsilon t} \sum_{i=1}^{n} \alpha_{i} |\Delta x_{i}(t,\phi,\varphi)|$$
  
$$\ge \min_{1 \le j \le n} \alpha_{j} e^{\varepsilon t} \sum_{i=1}^{n} |\Delta x_{i}(t,\phi,\varphi)|$$
  
$$\ge \min_{1 \le j \le n} \alpha_{j} e^{\varepsilon t} ||x_{i}(t,\phi) - x_{i}(t,\varphi)||_{\infty},$$
(7)

and

$$V(0) = \sum_{i=1}^{n} \alpha_{i} \bigg[ |x_{i}(0,\phi) - x_{i}(0,\varphi)| \\ + \sum_{j=1}^{n} \int_{-\tau_{j}(0)}^{0} \frac{L_{j}|c_{ij}|}{1 - \dot{\tau_{j}}(\psi_{j}^{-1}(s))} \\ \cdot \Delta x_{j}(s,\phi,\varphi) e^{\varepsilon(s+\tau_{j}(\psi_{j}^{-1}(s))} ds \bigg] \\ \leqslant \sum_{i=1}^{n} \alpha_{i} \bigg[ |\phi_{i}(0) - \varphi_{i}(0)| \\ + \sum_{j=1}^{n} \int_{-\tau}^{0} L_{j} \delta_{j}|c_{ij}| e^{\varepsilon\tau} |\phi_{j}(s) - \varphi_{j}(s)| e^{\varepsilon s} ds \\ \leqslant \sum_{i=1}^{n} \alpha_{i} \bigg[ 1 + \frac{L_{j} \delta_{j}}{\varepsilon} |c_{ij}| (e^{\varepsilon\tau} - 1) \bigg] \|\phi - \varphi\|_{\infty} (8)$$

This implies that the conclusion of the theorem hold by using Eq. (6)- (8).  $\Box$ 

**Proof of Theorem 1.** First, we prove that Eq. (1) possesses an  $\omega$ -periodic solution. For each solution  $x(t, \phi)$  of Eq. (1) and each  $t \ge 0$ , we can define a function  $x_t(\phi)$  in this fashion:

$$x_t(\phi)(\theta) = x(t+\theta,\phi), \text{ for } \theta \in [-\tau,0].$$

On this basis, we can define a mapping  $P: C^* \to C^*$  by

$$P\phi = x_{\omega}(\phi).$$

Let  $x(t, \phi)$ ,  $x(t, \varphi)$  be an arbitrary pair of solutions of Eq. (1). Let  $\varepsilon$  be a positive number satisfying Eq. (3). Let  $m \ge \frac{1}{\varepsilon\omega} \ln(2M(\varepsilon)) + 1$  be a positive integer. It follows from Lemma 3 that

$$P^{m}\phi - P^{m}\varphi \|$$

$$= \sup_{-\tau \leqslant \theta \leqslant 0} \|x(m\omega + \theta, \phi) - x(m\omega + \theta, \varphi)\|_{\infty}$$

$$\leqslant M(\varepsilon) \sup_{-\tau \leqslant \theta \leqslant 0} e^{-\varepsilon(m\omega + \theta)} \|\phi - \varphi\|_{\infty} ,$$

$$\leqslant M(\varepsilon) e^{-\varepsilon(m-1)\omega} \|\phi - \varphi\|_{\infty} \leqslant \frac{1}{2} \|\phi - \varphi\|_{\infty}$$

which shows that  $P^m$  is a contraction mapping on the Banach space  $C^*$ . According to the contraction mapping principle,  $P^m$  possesses a unique fixed point  $\phi^* \in C^*$ . Note that

$$P^m(P\phi^*) = P(P^m\phi^*) = P\phi^*$$

which indicates that  $P\phi^* \in C^*$  is also a fixed point of  $P^m$ . It follows from the uniqueness of fixed point of  $P^m$  that

ISBN: 978-988-19251-3-8 ISSN: 2078-0958 (Print); ISSN: 2078-0966 (Online) Proceedings of the World Congress on Engineering 2012 Vol I WCE 2012, July 4 - 6, 2012, London, U.K.

 $P\phi^* = \phi^*$ , viz.  $x_{\omega}(\phi^*) = \phi^*$ . Let  $x(t, \phi^*)$  be the solution of Eq. (1) with initial condition  $\phi^*$ , Then

$$x_{t+\omega}(\phi^*)(\theta) = x_t(x_{\omega}(\phi^*)) = x_t(\phi^*) \quad \text{for} \quad t \ge 0.$$

which implies

$$x(t+\omega,\phi^*) = x_{t+\omega}(\phi^*)(0) = x_t(x_{\omega}(0)) = x(t,\phi^*).$$

Thus,  $x(t, \phi^*)$  is  $\omega$ -periodic of Eq. (1).

On the other hand, it follows from Theorem 3 that every solution  $x(t,\phi)$  of Eq. (1) satisfies

$$\|x(t,\phi) - x(t,\phi^*)\|_{\infty} \leq M(\varepsilon) \|\phi - \phi^*\|_{\infty} e^{-\varepsilon}$$

for all  $t \ge 0$ . This shows that  $x(t, \phi)$  is globally exponentially periodic.  $\Box$ 

## III. AN ILLUSTRATIVE EXAMPLE

Consider the impulsive Hopfield neural network with timevarying delays:

$$\begin{pmatrix} \dot{x}_{1}(t) \\ \dot{x}_{2}(t) \end{pmatrix} = \begin{pmatrix} -2 & 0 \\ 0 & -2 \end{pmatrix} \begin{pmatrix} x_{1}(t) \\ x_{2}(t) \end{pmatrix} + \begin{pmatrix} 0.6 & 0.3 \\ 0.3 & -0.5 \end{pmatrix} \begin{pmatrix} \sin \frac{1}{\sqrt{2}} x_{1}(t) \\ \sin \frac{1}{2\sqrt{2}} x_{2}(t) \end{pmatrix} + \begin{pmatrix} 0.8 & -0.5 \\ -0.6 & 0.6 \end{pmatrix} \begin{pmatrix} \sin \frac{1}{\sqrt{2}} x_{1}(t - \tau_{1}(t)) \\ \sin \frac{1}{2\sqrt{2}} x_{2}(t - \tau_{2}(t)) \end{pmatrix}$$
(9)  
 +  $\begin{pmatrix} 1 - \cos 2\pi t \\ 1 + \sin 2\pi t \end{pmatrix},$   
  $\Delta x_{1}(t_{k}) = \gamma_{1k} x_{1}(t_{k}),$   
  $\Delta x_{2}(t_{k}) = \gamma_{2k} x_{2}(t_{k}).$ 

Obviously, the right hand side of Eq. (9) is 1-periodic(i.e.  $\omega = 1$ ). Now we investigate the influence of the delay and the period T of impulsive effect on the Eq. (9). If  $\tau(t) = \frac{1}{5}\pi$ , T = 1,  $\gamma_{1k} = \gamma_{2k} = 0.1$ , then p = 1 in (H<sub>2</sub>). According to Theorem 1, impulsive Hopfield neural networks Eq. (9) has a unique 1-periodic solution which is globally asymptotically stable(see Figs.1-Figs.4). In order to clearly observe the change trend of the solutions, we add a time coordinate axes to the Fig. 4 and change 2-D plan (Fig. 3) into 3-D space(Fig. 4).

If the effect of impulse is ignored, i.e.  $\gamma_{1k} = 0, \gamma_{2k} = 0$ , then Eq. (9) becomes periodic system. Obviously, the right hand side of Eq. (9) is 1-periodic. Numeric results show that Eq. (9) has a 1-periodic solution. Figs.5- Figs.8 show the dynamic behavior of the Eq. (9) with  $\tau(t) = \frac{1}{5}\pi$ . Furthermore, If  $\tau(t) = \frac{1}{5}\pi$  rises to  $\tau(t) = \pi$  gradually, then periodic oscillation of Eq. (9) will be destroyed. Numeric results show that Eq. (9) still has a global attractor which may be gui chaotic strange attractor(see Figs.9 -Figs.12). Every solutions of Eq. (9) will finally tend to the chaotic strange attractor.

## IV. CONCLUSION

We have established a sufficient condition for the existence and global exponential stability of a unique periodic solution in a class of HNNs with time-varying delays and periodic impulses, which assumes neither the differentiability nor the monotonicity of the activation functions.



Fig. 1. Time-series of the  $x_1(t)$  of Eq. (9) for  $t \in [0, 16]$ .



Fig. 2. Time-series of the  $x_2(t)$  of Eq. (9) for  $t \in [0, 16]$ .



Fig. 3. Phase portrait of 1-periodic solutions of Eq. (9) for  $t \in [0, 42]$ .



Fig. 4. pace figure of 1-periodic solutions of Eq. (9) by adding a time coordinate axes t.

Proceedings of the World Congress on Engineering 2012 Vol I WCE 2012, July 4 - 6, 2012, London, U.K.





Fig. 6. Time-series of the  $x_2(t)$  of Eq. (9) for  $t \in [0, 16]$  with  $\tau(t) = \frac{1}{5}\pi$ .



Fig. 7. Phase portrait of 1-periodic solutions of Eq. (9) with  $\tau(t) = \frac{1}{5}\pi$ .



Fig. 8. Space figure of 1-periodic solutions of Eq. (9) with  $\tau(t) = \frac{1}{5}\pi$ .







Fig. 10. Time-series of the  $x_2(t)$  of Eq. (9) for  $t \in [0, 48]$  with  $\tau(t) = \pi$ .



Fig. 11. Phase portrait of chaotic strange attractor of Eq. (9) with  $\tau(t) = \pi$ .



Fig. 12. Space figure of attractor of (9) by adding a coordinate axes t.

Proceedings of the World Congress on Engineering 2012 Vol I WCE 2012, July 4 - 6, 2012, London, U.K.

Our condition extends and generalizes a known condition for the global exponential periodicity of pure continuous Hopfield neural networks with time-varying delays. Further the numerical simulation shows that our system can occur many forms of complexities including chaotic strange attractor and periodic solution.

In recent years, numerous results have been reported on the stability of discrete as well as continuous neural networks. It is worthwhile to introduce various impulsive neural networks and then establish the corresponding stability results that include some known results for pure discrete or continuous neural networks as special cases.

#### REFERENCES

- [1] S. Arika and V. Tavsanoglu, "Global asymptotic stability analysis of bidirectional associative memory neural networks with constant time delays," Neurocomputing, vol. 68, pp. 161-176, Oct. 2005.
- J. Cao and Q. Jiang, "An analysis of periodic solutions of bi-[2] directional associative memory networks with time-varying delays," Physics Letters A, vol. 330, no. 3-4, pp. 203-213, Sept. 2004.
- [3] S. Li, X. Liao and C. Li, "Hopf bifurcation of a two-neuron network with different discrete time delays," Int. J. Bifurcation and Chaos, vol. 15, no. 5, pp. 1589-1601, 2005.
- [4] S. Xu, J. Lam and Y. Zou, "Delay-dependent approach to stabilization of time-delay chaotic systems via standard and delayed feedback controllers," Int. J. Bifurcation and Chaos, vol. 15, no. 4, pp. 1455-1465, 2005.
- [5] J. Liang and J. Lam, "Convergence of discrete-time recurrent neural networks with variable delay," Int. J. Bifurcation and Chaos, vol. 15, no. 2, pp. 581-595, 2005.
- [6] G. Chen, J. Zhou and Z. Liu, "Global synchronization of coupled delayed neural networks and applications to chaotic CNN models," Int. J. Bifurcation and Chaos, vol. 14, no. 7, pp. 2229-2240, 2004.
- [7] Z. Gui and W. Ge, "Impulsive effect of continuous-time neural networks under pure structural variations," *Int. J. Bifurcation and* Chaos, vol. 17, no. 6, pp. 2127-2139, 2007.
- Y. Li and C. Yang, "Global exponential stability analysis on impulsive [8] BAM neural networks with distributed delays," J. Math. Anal. Appl., vol. 324, no. 2, pp.1125-1139, Dec. 2006.
- X.-Y. Lou and B.-T. Cui, "Novel global stability criteria for high-order Hopfield-type neural networks with time-varying delays," J. Math. Anal. Appl., vol. 330, no. 1, pp. 144-158, June 2007.
- [10] X. Lou and B. Cui,"Delay-dependent stochastic stability of delayed Hopfield neural networks with Markovian jump parameters," J. Math. Anal. Appl., vol. 328,no. 1, pp. 316-326, Apr. 2007.
- [11] Y. Song, Y. Peng and J. Wei, "Bifurcations for a predator-prey system with two delays," J. Math. Anal. Appl., vol. 337, no. 1, pp.466-479, Jan. 2008.
- [12] S. Mohamad, "Global exponential stability in continuous-time and discrete-time delayed bidirectional neural networks," Physica D, vol. 159, no. 3-4, pp. 233-251, Nov. 2001.
- [13] X. Liao and C. Li, "An LMI approach to asymptotical stability of multi-delayed neural networks," *Physica D*, vol. 200, no.1-2, pp.139-155. Jan.2005.
- [14] J. Cao and L. Wang, "Exponential stability and periodic oscillatory solution in BAM networks with delays," IEEE Trans. Neural Networks, vol. 13, no. 2, pp. 456-463, Mar.2002.
- [15] X. Huang, J. Cao and D. Huang, "LMI-based approach for delaydependent exponential stability analysis of BAM neural networks,' Chaos, Solitons & Fractals, vol. 24, no. 3, pp. 885-898, May 2005.
- [16] D.-D. Bainov and P.-S. Simenov, Systems with Impulse Effect: Stability Theory and Applications, Ellis Horwood, Chichester, 1989.
- V. Lakshmikantham, D.-D. Baino and P.-S. Simeonov, Theory of Impulsive Differential Equations, World Scientific, Singapore, 1989.
- [18] K. Gopalsamy and B.-G. Zhang, "On delay differential equation with impulses," J. Math. Anal. Appl., vol. 139, no. 1, pp. 110C122, Apr.1989.
- [19] Z. Gui, X.-S. Yang and W. Ge, "Periodic solution for nonautonomous bidirectional associative memory neural networks with impulses," Neurocomputing, vol. 70, no. 13-15, pp. 2517-2527, Aug. 2007. [20] Z.-H. Guan and G. Chen, "On delayed impulsive Hopfield neural
- networks," Neural Networks, vol. 12, no. 2, pp.273C280, Mar. 1999.
- [21] H. Akca, R. Alassar, V. Covachev, Z. Covacheva and E. Al-Zahrani, "Continuous-time additive Hopfield-type neural networks with impulses," J. Math. Anal. Appl., vol. 290, no. 2, pp. 436-451, Feb. 2004.

- [22] M.-U. Akhmetov and A. Zafer, "Stability of the zero solution of impulsive differential equations by the Lyapunov second method," J. Math. Anal. Appl., vol. 248, no. 1, pp. 69C82, Aug. 2000.
- [23] Z. Gui, Biological Dynamic Models and Computer Simulation, Science Press, Beijing, 2005.
- [24] Z. Gui and W. Ge, "Existence and uniqueness of periodic solutions of nonautonomous cellular neural networks with impulses," Physics Letters A, vol.354, no. 1-2, pp. 84-94, May 2006.
- [25] Z. Gui and W. Ge, "Periodic solution and chaotic strange attractor for shunting inhibitory cellular neural networks with impulses," Chaos, vol. 16, no. 3 pp. 1-10, Aug. 2006. [26] Z. Gui and X.-S. Yang, "Stability and existence of periodic solutions of
- periodic cellular neural networks,"Computers and Mathematics With Application, vol. 52, no. 12, pp. 1657-1670, Dec.2006.
- [27] Z. Gui and D. Yuan, "Stability of artificial neural networks with impulsive inputs from outside the network," Journal of Computational Information Systems, vol. 2, no. 3, pp. 1045-1050, 2006.
- [28] Y. Li, "Global exponential stability of BAM neural networks with delays and impulses," Chaos, Solitons & Fractals, vol. 24, no. 1, pp. 279-285, Apr.2005.
- [29] Y. Li, W. Xing and L. Lu, "Existence and global exponential stability of periodic solution of a class of neural networks with impulses. Chaos, Solitons & Fractals, vol. 27, no. 2, pp. 437-445, Jan. 2006.
- [30] X. Yang, X. Liao, D.-J. Evans and Tang, Y. "Existence and stability of periodic solution in impulsive Hopfield neural networks with finite distributed delays," Physics Letters A, vol. 343, no. 1-3, pp.108-116, Aug. 2005.